

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:


- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/

HOW TO IDENTIFY * THE STARS *

QB 801 M5.











WILLIS-I-MILHAM




LIBRARY .


OF THE


University of California.

Class











THE MACMILLAN COMPANY NEW YORK · BOSTON · CHICAGO

NEW YORK · BOSTON · CHICAGO ATLANTA · SAN FRANCISCO

MACMILLAN & CO., LIMITED LONDON · BOMBAY · CALCUTTA MELBOURNE

THE MACMILLAN CO. OF CANADA, LTD. TORONTO

HOW TO IDENTIFY THE STARS

BY

WILLIS I. MILHAM, Ph.D.

FIELD MEMORIAL PROFESSOR OF ASTRONOMY
IN WILLIAMS COLLEGE



New York

THE MACMILLAN COMPANY

1909

All rights reserved

ME B801

GEREIM.

COPYRIGHT, 1909,

By THE MACMILLAN COMPANY.

Set up and electrotyped. Published June, 1909. Reprinted October, 1909.


Norwood Press

J. S. Cushing Co. — Berwick & Smith Co.
Norwood, Mass., U.S.A.

 \mathcal{N} .

CONTENTS

				PAGE
I.	Introduction	•	•	1
II.	THE HISTORY AND NUMBER OF THE COM	(S TEL)	LA-	
	TIONS	•	•	3
III.	THE METHODS OF DESIGNATING A STAR	•	•	12
IV.	STAR MAGNITUDES	•	•	15
v.	STAR COLORS	•	•	20
VI.	THE NUMBER OF THE STARS	•		21
VII.	THE METHOD OF LOCATING THE STA	RS A	ND	
	CONSTELLATIONS	•	•	23
VIII.	THE METHOD OF FURTHER STUDY .	•		33





I. INTRODUCTION

THE ability to recognize the more brilliant stars and to locate the more conspicuous constellations or star groups, is both an interesting and a useful acquirement. The number of people who have a real interest in popular astronomy and a fair acquaintance with the stars and constellations is steadily increasing and they find real pleasure in their information. the meteorologist who would make observations of the position of the streamers of the Aurora Borealis or locate the place of the appearance and disappearance of bright meteors, a knowledge of the stars and constellations is absolutely necessary. The best way to locate the position of an auroral streamer without the use of apparatus is to observe the stars between which and over which it extends. Similarly the point of appearance and disappearance of a meteor has been definitely determined if the nearest star in each case has been noted. If great exactness is desired, the angular distance and direction of the point in question from the nearest star may be estimated. Whenever an object is thus located by means of the stars, the exact time of observation and the place of observation (latitude, longitude, and elevation) must also be stated. From such observations made at two, preferably more, stations, the distance, height, and direction of an auroral display, and the distance, height, direction of motion, velocity of motion, and orbit in space of a meteor may be determined by computation.

The purpose of the constellation tracings and the descriptive material here given, is to serve as a guide in taking the first steps in learning the stars and constellations and also to point the way to the acquisition of further information on the part of those who desire it. The method here followed and the material presented is essentially the same as that used in the course on Descriptive Astronomy in Williams College.



II. THE HISTORY AND NUMBER OF THE CONSTELLATIONS

THE first definite information concerning the constellations comes from Claudius Ptolemy at the Alexandrian School of Philosophy in about 150 A.D. He was a great systematizer of information and summarized the astronomical learning of his time. His original writings in the Greek are lost but an Arabic translation of his manuscript on Astronomy, called the "Almagest," has come down to us. In it he enumerates, describes, and locates 48 constellations; 21 northern, 15 southern, and 12 zodiacal. These constellations by no means covered the whole sky, as a large part of the southern hemisphere was not mapped and there were often unmapped spaces between the constellations. As the centuries passed, the need of more constellations became apparent, but it was considered sacrilegious to add to the Ptolemaic list. Finally, in 1601, two constellations were added by Tycho Brahe. This opened the way, and for two centuries nearly every astronomer of note considered it his duty to map a group of stars and give to the group a name of his own choosing. Thus by 1800 there were at least 109 fairly well known constellations, and a large constellation (Argo) in the southern hemisphere was divided into four, thus making 112. The difficulties were now as great as before new constellations had been added, for some of the constellations overlapped and names had been given to some of them which could never become international. Gradually, by practically common consent, 24 were dropped, so that at present there are 88 constellations. The accompanying list shows the number of constellations added by each astronomer.

	HIS	TO	RY O	FI	HE CONS	TELL	ATIO	NS	5
140	A.D.		•		Ptolemy .	•	•	•	48
1601		•	•	•	Tycho Bra	he .	•	•	2
1603		•	•	•	Bayer .	•	•	•	12
1679		•	•		Royer .	•	•		5
1690		•	•		Halley .	•	•		1
1690		•	•	•	Hevelius .	•	•		11
1725		•	•	•	Flamsteed	•	•		2
1752		•		•	La Caille.		•	•	14
1770		•	•	•	Hell .	•	•		I
1776					Le Monnie	r .	•	•	2
1776		•	•	•	Lalande .	•	• .		I
1777		•			Poczobut	•	•		I
1800		•	•	•	Bode .	•	•	•	9
									109
Arg	go divi	ided	into f	our	parts .	•	•	•	3
Tw	enty-f	our	were d	lropp	ped	•	•	•	24 88

Some authorities attribute one constellation to the Emperor Hadrian in 130 A.D., one to Eratosthenes in 300 A.D., and four to Bartschius in 1624 A.D. and thus a smaller number to some mentioned in the table.

These eighty-eight constellations do not overlap, have definite boundaries, and cover the whole sky. The boundary lines usually wind in and out among the stars and are very irregular, particularly in the case of the

northern constellations. Some of the southern constellations have boundaries which are for considerable distances arcs of great or small circles. The constellations are also of very unequal size, some being ten times as large as others.

The following list contains the names of the constellations, the Latin genitive of the name, the meaning of the name, and the name of the originator of the constellation:—

ROPOSER
emy
Caille
er
emy
emy
emy
emy
emy .
emy
Caille
elius or artschius
emy
elius
eee



NAME GENITIVE MEANING PROPOSER Ca'-nis Ca'-nis Greater dog Ptolemy Majo'-ris Ma'-jor - Ca'-nis Ca'-nis Lesser dog **Ptolemy** Mi'-nor Mino'-ris - Capricor'-nus Capricor'-ni **Ptolemy** Goat Cari'-na Cari'-næ Keel La Caille -Cassiope'-ia Cassiopeia Ptolemy Cassione'-iæ Centau'-rus Centau'-ri Centaur Ptolemy - Ce'-pheus Ce'-phei Cepheus Ptolemy -Ce'-tus Ce'-ti Whale Ptolemy Chamæ'-leon Chamælon'-Chameleon Bayer tis Cir'-cinus Cir'-cini La Caille Compasses Colum'-ha Colum'-bæ Dove Royer or Bartschius Tycho Brahe -- Co'-ma Co'-mæ Berenice's Bereni'-ces Bereni'-ces hair or **Eratosthenes** Coro'-næ Coro'-na Southern Ptolemy Austra'-lis Austra'-lis crown Coro'-na Coro'-næ Northern **Ptolemy** Borea'-lis Borea'-lis crown Cor'-vus Cor'-vi Crow Ptolemy --Cra'-ter Crate'-ris **Ptolemy** Cup Crux Cru'-cis Cross Royer -Cyg'-nus Cyg'-ni **Ptolemy** Swan ~Delphi'-nus Delphi'-ni Dolphin Ptolemy Dora'-do Dora'-dus Sword-fish Bayer ' --Dra'-co Draco'-nis Dragon Ptolemy --- Equu'-leus Equu'-lei Little horse Ptolemy

NAME	GENITIVE	MEANING	Proposer
Erid'-anus	Erid'-ani	The river	Ptolemy
For'-nax	Forna'-cis	Furnace	La Caille
√Gem'-ini	Gemino'-rum	Twins	Ptolemy
Grus	Gru'-is	Crane	Bayer
Her'-cules	Her'-culis	Hercules	Ptolemy
Horolo'-gium	Horolo'-gii	Clock	La Caille
─Hy'-dra	Hy'-dræ	Snake	Ptolemy
Hy'-drus	Hy'-dri	Water snake	Bayer
In'-dus	In'-di	Indian	Bayer
Lacer'-ta	Lacer'-tæ	Lizard	Hevelius
-Le'-o	Leo'-nis	Lion	Ptolemy
Le'-o Mi'-nor	Leo'-nis Mino'-ris	Lesser lion	Hevelius
—T al mua	Lep'-oris	Hare	Dialoman
—Le'-pus —Li'-bra	Lep-ons Li'-bræ	Balance	Ptolemy
		Wolf	Ptolemy
Lu'-pus	Lu'-pi		Ptolemy
Lynx	Lyn'-cis	Lynx	Hevelius
_Ly'-ra	Ly'-ræ	Harp	Ptolemy
Ma'-lus	Ma'-li	Mast	La Caille
Men'-sa	Men'-sæ `	Table	La Caille
Microsco'- pium	Microsco'-pii	Microscope	La Caille
Monoc'-eros	Monocero'-tis	Unicorn	Hevelius or Bartschius
Mus'-ca	Mus'-cæ	Fly	Bayer
Nor'-ma	Nor'-mæ	Rule	La Caille
Oc'-tans	Octan'-tis	Octant	La Caille
Ophiu'-chus	Ophiu'-chi	Serpent carrier	Ptolemy
Ori'-on	Orio'-nis	Orion	Ptolemy

NAME	GENITIVE	MEANING	Proposer
Pa'-vo	Pavo'-nis	Peacock	Bayer
Peg'-asus	Peg'-asi	Winged horse	Ptolemy
→Per'-seus	Per'-sei	Perseus	Ptolemy
Phœ'-nix	Phœni'-cis	Phœnix	Bayer
Pic'-tor	Picto'-ris	Painter	La Caille
Pis'-ces	Pis'-cium	Fishes	Ptolemy
Pis'-cis	Pis'-cis	Southern	Ptolemy
Austra'-lis	Austra'-lis	fish	
Pup'-pis	Pup'-pis	Stern	La Caille
Retic'-ulum	Retic'-uli	Net	La Caille
Sagit'-ta	Sagit'-tæ	Arrow	Ptolemy
Sagitta'-rius	Sagitta'-rii	Archer	Ptolemy
Scor'-pius	Scor'-pii	Scorpion	Ptolemy
Sculp'-tor	Sculpto'-ris	Sculptor	La Caille
Scu'-tum	Scu'-ti	Shield	Hevelius
Ser'-pens	Serpen'-tis	Serpent	Ptolemy
Sex'-tans	Sextan'-tis	Sextant	Hevelius
Tau'-rus	Tau'-ri	Bull	Ptolemy
Telesco'-pium	Telesco'-pii	Telescope	La Caille
Trian'-gulum	Trian'-guli	Triangle	Ptolemy
Trian'-gulum	Trian'-guli	Southern	Bayer
Austra'-le	Austra'-lis	triangle	
Tuca'-na	Tuca'-næ	American goosa	Bayer
-Ur'-sa	Ur'-sæ	Greater bear	D4 - 1
Ma'-jor	Majo'-ris	Greater bear	Ptolemy
⊸Ur'-sa	Ur'-sæ	Lesser bear	Ptolemy
Mi'-nor	Mino'-ris		,
Ve'-la	Velo'-rum	Sails	La Caille
√Vir'-go	Vir'-ginis	Virgin	Ptolemy
Vo'-lans	Volan'-tis	Flying (fish)	Bayer
Vulpec'-ula	Vulpec'-ulæ	Fox	Hevelius

It will be seen that there are three kinds of names used; names of animals, names borrowed from mythology, and names of pieces of apparatus.

When and where the constellations were originally designed are questions which have received a great deal of attention and study. This subject can be merely sketched in outline here. The list of 48 constellations given by Ptolemy was by no means original with him, but was simply a slight revision of an earlier list by Hipparchus, date about 140 B.C. And even before his time a fairly full description of the constellations can be found in the poem of Aratus of Soli, date about 280 B.C. But this was simply a versification of an earlier astronomical work by Eudoxus in 370 B.C., and this in turn, as we now know, was not based on observations made by the Greeks, but on astronomical information that was at that time at least two thousand years old. There are three sources of information as to the origin of the constellations: first, the internal evidence from the constellations themselves; secondly, the documentary evidence in the form of references in the early writings; thirdly, the inscriptions on monuments, tablets, coins, etc., which are continually being discovered. The originators of the constellations naturally portrayed the things with which they were familiar. The elephant, camel, tiger, and crocodile are not represented. This would exclude India and Egypt as possible places of origin. The portion of the sky left unmapped shows the portion of the sky not visible to them. They must thus have lived in about 38° north latitude. In short, all lines of evidence show that the constellations probably originated in the Euphrates valley, and were practically complete as early as 3000 B.C. What we have is thus the Latin form of a Greek digest of very early information. For further information about the origin of the constellations, see—

BROWN, Primitive Constellations; Williams and Norgate, London, 1899.

LOCKYER, The Dawn of Astronomy; The Macmillan Company, 1894.

MAUNDER, Astronomy without a Telescope, Chapter I; "Knowledge" Office, London, 1902.

MAUNDER, The Oldest Picture-book of All; The Nineteenth Century Magazine, September, 1900.

III. THE METHODS OF DESIGNATING A STAR

The oldest, but now obsolete, method of designating a particular star was to describe its position in the constellation. The portion of the sky belonging to a certain constellation was supposed to be covered by a representation or picture of the thing after which the constellation was named. Thus a star might be described as located "in the head of Hercules" or "in the right knee of Bootes" or "in the horn of the bull." Many star charts still give those old pictures in connection with the constellations.

There are four modern methods of designating a star. (1) By name. Hundreds of stars have received individual names, but less than a hundred of these are now in general use. They are usually of Latin, Greek, or Arabic origin, and are either proper names or describe position. For example: Regulus, Spica,

Aldebaran. (2) By means of a letter. In 1603 Bayer introduced the system of designating the stars in a constellation by means of the letters in the Greek alphabet. These are, in order:—

α alpha	η eta	ν nu	au tau
$\boldsymbol{\beta}$ beta	heta theta	ξ xi	v upsilon
γ gamma	ιiota	o omicron	ϕ phi
δ delta	κ kappa	π pi	χ chi
ϵ epsilon	λ lambda	ho rho	ψ psi
ζ zeta	μ mu	σ sigma	ω omega

The stars were usually lettered in the order of brightness, a being the brightest star, β the next brightest, and so on. When the letters of the Greek alphabet were exhausted, the Roman letters were used. In a few instances the stars were lettered in order of position instead of brightness. The Latin genitive of the constellation name always follows the Greek or Roman letter. For example: a Lyræ, β Geminorum, g Ursæ Majoris. (3) By number. About 1700 Flamsteed introduced the custom of numbering the stars in a constellation. This applies usually to the

fainter stars which had not been lettered, and they were numbered in order of position from west to east. For example: 61 Cygni, 50 Cassiopeiæ. (4) By catalogue number. In the case of some faint stars visible to the naked eye, and nearly all stars visible in a telescope only, the star is designated as having a certain number in a certain catalogue. It is usually the first catalogue in which it occurred or some very well known comprehensive catalogue. For example: Groombridge 966, B.A.C. 4536.

The last three of the four modern methods seldom overlap at present. The brighter stars are usually lettered. Those which are fainter, but still visible to the naked eye, are usually numbered, while telescopic stars are referred to simply by catalogue and number. If a star has a proper name, this is usually given in addition to the Bayer letter.

IV: STAR MAGNITUDES

THE term "magnitude" when applied to a star refers simply to its brightness and not to its volume or mass. Ptolemy arbitrarily graded the stars visible to the naked eye into six classes or magnitudes, the sixth being the faintest and the first the brightest. The reason for having six magnitudes rather than more is not known, unless it is because the eye without the help of instruments is unable to make finer distinctions. After the telescope was invented, the number of magnitudes had to be increased so as to include these faint telescopic stars, but there was no uniformity among different observers in extending the system. As the result, the magnitude of a faint star, as determined by different observers, might differ by even two or three. As measurements became more exact. it also became desirable to subdivide magnitudes.

It had been noticed for some time that, roughly speaking, a star of the first magnitude was about one hundred times as bright as a star of the sixth. In 1850, Pogson proposed to make the $\sqrt[5]{100}$, or 2.512, the uniform ratio between successive magnitudes.

Thus, $b_1 = \sqrt[5]{100} \ b_2$, where b_1 is the brightness of a first magnitude star and b_2 that of a second.

Similarly,
$$b_2 = \sqrt[5]{100} \ b_3,$$

$$b_3 = \sqrt[5]{100} \ b_4 \text{ etc.}$$
From these
$$b_1 = (\sqrt[5]{100})^2 b_3,$$
or
$$b_1 = (\sqrt[5]{100})^4 b_5,$$
or
$$b_3 = (\sqrt[5]{100})^6 b_9.$$
Thus, in general,
$$b_m = (\sqrt[5]{100})^{n-m} b_n.$$
Transposing this,
$$\frac{b_m}{b_n} = (\sqrt[5]{100})^{n-m},$$

$$\log_{10} \frac{b_m}{b_n} = (n-m) \log_{10} \sqrt[5]{100},$$

$$\log_{10} \frac{b_m}{b_n} = 0.4 (n-m).$$

This formula put into words is: the logarithm of the ratio of brightness equals four tenths of

the difference in magnitude. By means of this formula, if the magnitudes of two stars are given, the ratio of brightness may be computed, and if the ratio of brightness and the magnitude of one star is given, the magnitude of the other may be computed.

EXAMPLE 1. If the magnitude of Mizar (ζ Ursæ Majoris) is 2.4 and the magnitude of Alkor (g Ursæ Majoris) is 4.0, find their ratio of brightness.

$$log_{10}$$
 ratio = 0.4(4.0 - 2.4) = 0.4(1.6) = 0.64.
Ratio of brightness = 4.37.

Example 2. If a star is 30 times as bright as Polaris (a Ursæ Minoris), magnitude 2.1, find its magnitude.

$$\log_{10}30 = 0.4(2.1 - x),$$

$$1.477 = 0.84 - 0.4 x,$$

$$0.4 x = -0.637,$$

$$x = -1.59.$$

The magnitude is -1.59, and this is the magnitude of Sirius (a Canis Majoris).

This definite system of star magnitudes was gradually adopted, and now all star magnitudes are expressed in it. Some star must be considered to be fundamental, that is, to have a definite unchanging magnitude, and the magnitudes of the other stars must be determined in terms of its magnitude. Polaris (a Ursæ Minoris) is usually taken as the fundamental star. By means of photometers all other stars are compared with it, either directly or indirectly, and the ratio of brightness, and thus the magnitude, determined. The work of the Harvard College Observatory in this connection has been particularly noteworthy. Here the actual standard is not Polaris but the mean magnitude of one hundred circumpolar stars of about the fifth magnitude.

In this system of magnitudes, the naked eye can perceive stars of about the sixth magnitude if the air is clear and free from haze. One disadvantage of the system is the fact that there are stars brighter than the first magnitude. A star 2.51 times as bright as one of the first magnitude would be of the o magnitude. A star 2.51 times as bright as that would be of the —1 magnitude. Thus for a very bright object a negative magnitude must be used. There is one star which has

a negative magnitude, Sirius (a Canis Majoris) -1.6. On the same scale the sun would have a magnitude of -26.3.

The magnitudes of stars remain remarkably constant. There are only about 500 stars in connection with which a variation in magnitude has been detected, and only about 50 of these are visible to the naked eye. The change in magnitude may be a steady change in one direction, an irregular fluctuation, or a periodic variation. If the fluctuation is irregular, only the limits can be stated. If the variation is periodic, such facts as the limits of the variation and the period may be determined.

V. STAR COLORS

Most of the stars appear to us as white, although in a few cases the star appears to be tinged with color. The following list probably includes the various colors as they appear to the naked eye: greenish white, bluish white, white, yellowish white, orange-white, reddish white. Stars are sometimes said to be yellow, orange, or red, but in every case the white predominates to such an extent that it is really only a tinge of the color in question that is perceived.

As seen through the telescope all the spectrum colors may be noticed, but in every case they are strongly mixed with white.

The causes of these differences in color are differences in temperature and chemical composition.

VI. THE NUMBER OF THE STARS

THE opinion is often expressed that the stars appear almost countless in number. As a matter of fact the total number visible to the naked eye is only about 6,000. Only one-half of these are visible at any one time, and a small amount of haze greatly reduces the number that is visible, particularly near the horizon. Thus on a night which would be considered clear only about 2,000 stars are visible. As was said before, the sixth magnitude is usually considered the limit of visibility to the naked eye. The following table gives the number of stars between each magnitude. The second line of the table gives the values of 4×3^m , where m is the magnitude. The agreement is remarkably close. Expressed in words, the law would be that there are about three times as many stars in each magnitude as in the preceding magnitude.

Magnitude o	1	2	3	4	5	6
Number of stars	11	28	103	321	1020	2843
4×3 ^m	I 2	36	108	324	972	2916

The following list of the twenty brightest stars gives the proper name, the Bayer letter, the magnitude, and the color. The word "(invisible)" indicates that the star is never visible in the United States:—

THE TWENTY BRIGHTEST STARS

Sirius	a Canis Majoris -	- 1 . 6	Bluish white
Vega	a Lyræ	0.1	Bluish white
Arcturus	a Bootis	0.2	Orange
Capella	a Aurigæ	0.2	Yellow
Rigel	β Orionis	0.3	White
	a Centauri	0.3	White (invisible)
Canopus	a Argus	0.4	Bluish white (invisible)
Procyon	a Canis Minoris	0.5	White
Achernar	a Eridani	0.6	White (invisible)
Betelgeuse	a Orionis	0.9	Red
Altair	a Aquilæ	0.9	Yellow
	β Centauri	0.9	White (invisible)
Aldebaran	a Tauri	1.1	Red
Antares	a Scorpii	1.2	Red
Pollux	β Geminorum	1.2	Orange
Spica	a Virginis	1.2	White
	a Crucis	1.3	Bluish white (invisible)
Fomalhaut	a Piscis Australis	1.3	Orange
Regulus	a Leonis	1.3	White
Deneb	a Cygni	1.3	White

VII. THE METHOD OF LOCATING THE STARS AND CONSTELLATIONS

THE method of locating the stars and constellations here advocated is by means of the so-called constellation tracings. Most of the conspicuous constellations contain bright stars which, when connected by lines, form figures, such as a triangle, square, W, sickle, etc., which are easily remembered and quickly recognized. After these tracings have once been learned and noticed, as soon as the eye falls upon that part of the sky, it instinctively runs through the tracing, and the constellation is thus recognized and its stars identified. For convenience in locating them, the 88 constellations have been divided into four groups. The first group consists of 28 conspicuous constellations with tracings. The second group contains the five constellations which are chiefly conspicuous on account of a single very bright star. The third group of 19 comprises the

HOW TO IDENTIFY THE STARS

24

inconspicuous constellations, while the fourth group contains the 36 constellations which are too far south to be seen at all or well seen from the United States. The following table in four parts gives these four groups of constellations:—

THE 88 CONSTELLATIONS DIVIDED INTO FOUR GROUPS

I. THE 28 CONSPICUOUS CONSTELLATIONS WITH TRACINGS

Andromeda		LYRA (triangle and parallelogram)		
AQUILA		OPHIUCHUS		
ARIES	(triangle)	ORION	(parallelogram)	
Auriga	(pentagon)	PEGASUS	(square)	
CASSIOPEIA	(W)	Perseus		
CORONA BOREALI	S (semicircle)	SAGITTA	(arrow)	
Corvus		SAGITTARIUS		
Cygnus	(cross)	Scorpius	(sickie)	
DELPHINUS		SERPENS		
Draco		TAURUS	(wedge)	
Gemini		TRIANGULUM	(triangle)	
Hercules		Ursa Major	(big dipper)	
LEO (sickle	and triangle)	URSA MINOR	(little dipper)	
Libra		Virgo		

II. THE 5 CONSTELLATIONS WITH A SINGLE VERY BRIGHT STAR

BOOTES CANIS MAJOR
CANES VENATICI CANIS MINOR
PISCIS AUSTRALIS

III. THE 19 INCONSPICUOUS CONSTELLATIONS

AQUARIUS ERIDANUS
CAMELOPARDALIS HYDRA
CANCER LACERTA
CAPRICORNUS LEO MINOR
CEPHEUS LYNX

CETUS MONOCEROS
COMA BERENICES PISCES
CRATER SCUTUM

CRATER SCUTUM
EQUULEUS SEXTANS

Vulpecula

IV. THE 36 SOUTHERN CONSTELLATIONS

Antlia Lupus
Apus Malus
Ara Mensa

Cælum Microscopium

MUSCA CARINA Norma CENTAURUS CHAMÆLEON **OCTANS** CIRCINUS Pavo COLUMBA PHŒNIX CORONA AUSTRALIS PICTOR CRUX Puppis DORADO RETICULUM **FORNAX** SCULPTOR GRUS TELESCOPIUM . Horologium TRIANGULUM AUS.

HYDRUS TUCANA
INDUS VELA
LEPUS VOLANS

Charts I. to IV. show the position of the constellations at 9 P.M. during January, April, July, and October. During a following month they would show the position two hours earlier, and during a preceding month two hours later. That is, stars come into the same position two hours earlier after the lapse of a month. The 28 conspicuous constellations and the five with single very bright stars are printed in small capitals, while the inconspicuous constellations are printed in small letters. In order to match the sky, the chart is supposed to be held over the head of an observer facing south. Thus for a convenient hour of the evening, the position of the constellations is given for every month in the year. In order to get the position at any other hour of the evening, only the apparent diurnal rotation of the heavens need be taken account of.

Figures 1 to 24 give the tracings and information about the 28 conspicuous constellations. Andromeda and Perseus, Aries and Triangulum, Libra and Scorpius, Ophiuchus and Serpens, are represented by one figure for each pair. The faint lines represent the tracings.

The arrow points to the north pole of the heavens, and thus indicates for a constellation in any position how the tracing must be held to match the sky. Three symbols are used to represent the stars, depending upon their magnitude: + for stars having a magnitude of 1.4 or brighter; • for stars having a magnitude between 1.5 and 3.4 inclusive; • for stars having a magnitude of 3.5 or less. The designation of the star and its exact magnitude according to the measurements made at the Harvard College Observatory are stated. The proper name of the star is also given, and its color if not white, and, if variable, the limits of the variation and the period if periodic. At the bottom of each figure the right ascension and declination of the center of each constellation are indicated.1

¹ Right ascension and declination are the coördinates in the so-called equator system of coördinates. The fundamental points of this system are the poles of the celestial sphere, that is, the points of no diurnal motion on the celestial sphere. These are the points where the earth's axis produced cuts the celestial sphere. The north pole is located near Polaris (a Ursæ Minoris). The celestial equator is a great circle 90° from the pole. It is the great circle in which the plane of the earth's equator cuts the celestial sphere. The equator runs through the con-

The facts concerning the five constellations with a single very bright star are:—

Bootes — α ; name, Arcturus; magnitude, 0.2; color, orange-white.

Canes Venatici—a; name, Cor Caroli; magnitude, 3.3; color, white.

Canis Major— α ; name, Sirius; magnitude, —1.6; color, bluish white.

Canis Minor—a; name, Procyon; magnitude, 0.5; color, bluish white. (β Canis Minoris has a magnitude of 3.1 and is fairly conspicuous. The constellation could perhaps have been represented by these two stars better than by a single star.)

Piscis Australis—a; name, Fomalhaut; magnitude, 1.3; color, orange-white.

Of the 19 inconspicuous constellations Cancer and Cepheus are perhaps the most easily recognized. There are two stars in Cepheus

stellations of Pisces, Cetus, Taurus, Orion, Monoceros, Hydra, Sextans, Leo, Virgo, Serpens, Ophiuchus, Aquila, Aquarius. The great circles drawn through the poles and perpendicular to the equator are called hour circles. The starting point or origin of coördinates is the vernal equinox, the point where the sun crosses the equator on March 21. It is located in the constellation of Pisces. Declination is angular distance north or south of the equator; plus when north and minus when south. Right ascension is the portion of the equator between the vernal equinox and the hour circle through the point in question. It is usually expressed in time and runs from 0 to 24 hours.

which catch the eye first and they are quite a little brighter than those near them. These are α (magnitude 2.6) and β (magnitude 3.3). Cancer is noticeable on account of the three stars quite close together which form a little equilateral triangle.

The best method of building up an acquaintance with the stars and constellations from the material here given can only be suggested. Most people know at least two or three constellations. If that is the case, the best method of procedure is to study first the constellations which surround those which are already known. Notice the tracing, determine where the constellation is with reference to those already known, and then try to pick it out among the stars. In this way all of the constellations will become gradually known. If no constellations at all are already familiar to the observer, start with those which are by far the most conspicuous. These are probably Aries, Auriga, Cassiopeia, Leo, Lyra, Orion, Pegasus, Scorpius, Taurus, Ursa Major. Determine from the charts where the constellation is located. observe carefully the tracing, noting particularly the magnitudes of the stars, and then persevere until the constellation is recognized.

For example, if the attempt is to be made to find the constellation of Taurus, fix firmly in mind before going out to observe the sky that the tracing is a V, that all of the stars but one are of about the same magnitude and only moderately bright, and that the one bright star is of the first magnitude and reddish in color. As these facts are sometimes forgotten while observing the sky, it is often convenient to take with one the tracings and a pocket electric lamp for illuminating them.

When a beginning has once been made, progress is usually easy. A single hour's instruction by one familiar with the stars and constellations is sufficient to give a considerable amount of information to a beginner, particularly if the charts and tracings have been carefully studied previously.

Another method advocated by some for building up an acquaintance with the stars and constellations is to start with the fifteen brightest stars. The chief facts about these stars have been given in a previous table. Find out, by looking up the constellations on the charts, how many of these are visible at the desired time of observation and in what part of the sky they are located. Then persevere as before until these have been recognized, and from these as known points build up an acquaintance with the surrounding stars and constellations. This method of starting with the bright stars can be used to the best advantage when the moon is from three to seven days old. The light of the moon pales out the fainter stars, thus making the bright ones more conspicuous. A perfectly clear moonless night is not the best time to begin observations, as the number of stars visible is confusing.

Some advocate trying to locate the stars and constellations by identifying them when rising or setting. The disadvantages of this method are that the horizon is always more hazy and cloud-covered than the rest of the sky, and, secondly, that it confines the time of observation to a particular hour of the night and is impossible at some times of year.

Any attempt to locate a star or constellation by giving its direction and distance from a known point will be no more successful than to notice its location on a chart.

VIII. THE METHOD OF FURTHER STUDY

THE charts, figures, and information given above can serve only as a guide in taking the first steps in learning the stars and constellations. Further knowledge is both useful and pleasant, and to gain it the reader must be referred to the numerous books bearing on the subject. Some of these books will be indicated here.

ALLEN, Star-names and their Meanings; G. E. Stechert, 1899, is a veritable mine of information. It gives the meaning of the names used in connection with the stars and constellations, numerous extracts from the writings where these names were used, and the more important facts such as magnitude, color, etc., about the chief stars in each constellation. It is a compendium of information, but not a book to be read through as a whole.

There are two books, MARTIN, The Friendly Stars; Harper and Brothers, 1907,

D

and SERVISS, Astronomy with the Naked Eye; Harper and Brothers, 1908, which attempt to put in readable and attractive form the facts concerning the stars and constellations. The last-named book also contains star charts, giving the position of all stars down to the sixth magnitude. These are the two books to recommend whenever a book giving a popular presentation of the subject is desired.

Among the modern up-to-date star atlases may be mentioned:—

BALL, A Popular Guide to the Heavens; George Philip & Son, London, 1905.

KLEIN, Star Atlas; E. & J. B. Young & Co., New York, 1901.

PECK, The Observer's Atlas of the Heavens; Gall & Inglis, London, 1898.

UPTON, Star Atlas; Ginn & Company, 1896.

MESSER, Stern-Atlas für Himmelsbeobachtungen; K. L. Ricker, Leipzig, 1902.

A star atlas is absolutely essential to one who desires to add to his knowledge of the stars and constellations. The star charts in these atlases represent all the constellations with their boundaries, and give the location of all stars visible to the naked eye. In addition

much information is given concerning star magnitudes, double stars, variable stars, star colors, etc. The meaning of the names, however, and the historical side of the subject are usually not treated. If it is desired to identify an unknown star, the best method is to estimate its magnitude carefully, and note its location with reference to several known stars or constellations. A chart of this portion of the sky may then be consulted, and the unknown star can usually be readily identified and the constellation to which it belongs determined. If it is desired to locate an inconspicuous star or constellation, the best method is to note on the appropriate chart its location with reference to several easily identified stars or constellations. On turning to the sky the inconspicuous object can usually be readily located and identified. In this way, by means of star atlases, one's information may be indefinitely extended.

Such atlases as ARGELANDER'S Atlas des Nördlichen Gestirnten Himmels (2d edition by Küstner, Bonn, 1899), PETERS'S Celestial Charts (published at Canton, N. Y.),

and the *Uranometria Argentina* are useful only in connection with the telescope when it is desired to identify all the stars, perhaps even to the ninth or tenth magnitude, within a very small area such as the field of view of the telescope.

Lists of the constellations, the stars visible to the naked eye, colored stars, variable stars, etc., can also be found in CHAMBERS'S Handbook of Descriptive and Practical Astronomy (The Clarendon Press, 1890) and VALENTINER, Handwörterbuch der Astronomie (Breslau, 1901).

If the coördinates of a star, right ascension, and declination are desired, they may be found for a small selected list of stars in *The American Ephemeris and Nautical Almanac*, published each year at Washington, or in the corresponding year books of the English, French, and German governments. AMBRONN, *Sternverzeichnis* (Julius Springer, Berlin, 1907) also contains the coördinates of all stars (7796 in all) which are above the 6.5 star magnitude. Such information, however, is desired not by the observer who is locating an object by

¢

means of the stars, but by the computer who is reducing the observations.

The following list contains other valuable books on the stars and constellations:—

BURRITT, Atlas designed to illustrate the Geography of the Heavens; New York, 1835.

COTTAM, Charts of the Constellations; London, 1889. DIEN, Atlas Céleste; Paris, 1869.

GORE, Star Groups; Crosby Lockwood & Sons, London, 1891.

HEIS, Atlas Cælestis Nouvus; Cologne, 1872.

HILL, The Stars and Constellations; Funk & Wagnalls Company, New York, 1894.

JEANS, Handbook for Finding the Stars; London, 1888. JOHNSTON, School Atlas of Astronomy (revised by Grant); G. P. Putnam's Sons, New York.

MAUNDER, Astronomy without a Telescope; London, 1902.

PECK, The Constellations and how to find them; Gall & Inglis, London.

PORTER, The Stars in Song and Legend; Ginn & Company, New York, 1902.

PROCTOR, Half-hours with the Stars; G. P. Putnam's Sons, 1875.

SERVISS, *Pleasures of the Telescope*; D. Appleton & Company, New York, 1905.

SERVISS, Astronomy with an Opera-glass; D. Appleton & Company, New York, 1906.

YOUNG, Uranography, in his Elements of Astronomy; New York, 1890. The star atlases of Burritt, Heis, and Dien are particularly interesting, as they give great prominence to the constellation figures, as was the case with all the older atlases.



CHART I.







CHART II.






CHART III.







CHART IV.



•




Fig. 1




Fig. 2






FIG. 3




Fig. 4






FIG. 5





Fig. 6



ŧ ,

.

.′

•




Fig. 7





FIG. 8





ļ

FIG. 9



FIG. 10

÷




Fig. 11





FIG. 12





١

FIG. 13




FIG. 14






FIG. 15




FIG. 16






FIG. 17




FIG. 18






FIG. 19




FIG. 20

•




FIG. 21




FIG. 22






FIG. 23




FIG. 24





BOOKS FOR THE GENERAL READER OF SCIENTIFIC TASTES

Avebury's Peace and Happiness

By the Right Honorable Lord Avebury, P.C. (formerly Sir John Lubbock). 12mo. Cloth. 306 pages. \$1.50 net.

Simple, sensible advice on such topics as the proper treatment of the body and the mind, on education, on our attitude towards children, friends, and enemies, on riches, on nature and religion.

Bailey's The Evolution of Our Native Fruits

By L. H. Bailey. 12mo, Cloth, Illus. 472 pages, \$2.00 net.

In this entertaining volume, the origin and development of the fruits peculiar to North America are inquired into, as well as the personality of those horticultural pioneers whose almost forgotten labors have given us our most valuable fruits.

Bailey's The State and the Farmer

By L. H. Bailey. 12mo. Cloth. 189 pages. \$1.25 net.

A book discussing the political and economic status of the farmer. Of particular interest not only to the agriculturist but also to the student of economics and to the general reader.

Bailey's The Survival of the Unlike

A Collection of Evolution Essays suggested by the Study of Domestic Plants. By L. H. Bailey. 515 pages. \$2.00 net.

This work treats the modification of plants under cultivation, and incidentally gives much that is suggestive and valuable to the working horticulturist,

Ely's A Woman's Hardy Garden

By Helena Rutherfurd Ely. With illustrations from photographs. 12mo, Cloth. 231 pages. \$1.75 net.

Elv's Another Hardy Garden Book

By Helena Rutherfurd Ely. With illustrations from photographs. 12mo, Cloth. 258 pages. \$1.75 net.

"Mrs. Ely is the wisest and most winning teacher of the fascinating art of gardening that we have met in modern print."—New York Tribune.

French's A Book of Vegetables and Garden Herbs

A Practical Hand-book and Planting Table for the Home Garden. By Allen French. 12mo. Cloth. Illus. 338 pages. \$1.75 net.

Hall's Three Acres and Liberty

By Bolton Hall. 12mo. Cloth. Illus. 457 pages. \$1.75 net.

Possibilities of the small suburban farm, and practical suggestions to city dwellers how to acquire and make profitable use of it.

Harwood's New Creations in Plant Life

By W. S. Harwood. 12mo. Cloth. Illus. 439 pages. \$1.75 net.

This is the first authentic account of the really wonderful experiments with plant life conducted by Mr. Luther Burbank in California.

Harwood's The New Earth

A Recital of the Triumphs of Modern Agriculture in America. By W. S. Harwood. 12mo. Cloth. Illus. 390 pages. \$1.75 net.

Hutchinson's The Story of the Hills

A Book about Mountains for General Readers, By Rev, H, W, Hutchinson, 12mo. Cloth. Illus. 372 pages. \$1.50.

"A clear account of the geological formation of mountains and their various methods of origin in language so clear and untechnical that it will not confuse even the most unscientific."—Boston Evening Transcript.

Illinois Girl's A Prairie Winter

By an Illinois Girl. Cloth, 164 pages, \$1.00 net.

A record of the procession of the months, from midway in September to midway in May.

Ingersoll's The Life of Animals: The Mammals

By Ernest Ingersoll, With Fourteen Full-page Color Plates and Many Other Illustrations. 8vo. Cloth. 566 pages. \$2.00 net.

"Each of the groups of mammals from the highest to the lowest is taken up in turn with illustrations, descriptions, and anecdotes of some of the representatives. Many of the illustrations are from original photographs and drawings, and greatly enhance the value of the work."

- American Journal of Science, New Haven.

Ingersoll's Wild Neighbors

Outdoor Studies in the United States, By Ernest Ingersoll, With Numerous Illustrations, 12mo, Cloth, 313 pages, \$1.50.

Studies and stories of the gray squirrel, the puma, the coyote, the badger and other burrowers, the porcupine, the skunk, the woodchuck, and the raccoon.

McFarland's Getting Acquainted with the Trees

By J. Horace McFarland. &vo. Cloth. Illus. 150 pages. \$1.50 net.

Marble's Nature Pictures by American Poets

Selected and Edited by Annie Russell Marble, 12mo, Cloth, 248 pages, 50 cents net.

Mead's Irrigation Institutions

A Discussion of the Economic and Legal Questions created by the Growth of Irrigated Agriculture in the West. By Elwood Mead. 12mo. Half leather. 403 pages. \$1.25 net.

This book is based upon twenty years' experience in the development of irrigated agriculture in the West under conditions, official and otherwise, which gave unprecedented opportunities for information and judgment.

Peterson's How to Know the Wild Fruits

A Guide to Plants when not in Flower, by Means of Fruit and Leaf. By Maude Gridley Peterson. Illustrated by Mary Elizabeth Herbert. 12mo. Cloth. 482 pages. \$1.50 net.

Scott's The Story of a Bird Lover

By William Earl Dodge Scott, 12mo, Cloth, Illus, 383 pages, \$1.50 net.

"There is charm in the manner of the telling of this bird-lover story which makes it appeal not alone to him who has scientific names at his tongue's tip, and who knows the nesting day of every bird in the ornithological list, but to him as well who has simply the layman's love for a breezy narrative and for that freshness which comes from a well-told tale of the free fields."

- Chicago Evening Post.

Sheldon's The Farm and the Dairy

By J. P. Sheldon. 12mo. Cloth. Illus. 166 pages. \$1.00 net.

A valuable text-book on practical dairying, detailing the most modern developments in dairy farming.

Smythe's The Conquest of Arid America

By William E. Smythe. 8vo. Cloth. Illus. 360 pages. \$1.50 net.

Mr. Smythe discusses colonization and irrigation in a general way, and the intervention of the Government in the task of reclaiming the desert parts of our country.

St. Maur's A Self-Supporting Home

By Kate V. St. Maur. 12mo. Cloth. Illus. 356 pages. \$1.75 net.

Each chapter is the detailed account of all the work necessary for one month—in the vegetable garden, among the small fruits, with the fowls, guineas, rabbits, cavies, and in every branch of husbandry to be met with on the small farm.

"One of the most sensible, practical books of the kind ever published."

- Louisville Courier-Journal.

St. Maur's The Earth's Bounty

By Kate V. St. Maur. 438 pages. \$1.75 net.

Written in the same practical, sensible, and thoroughly interesting vein as the preceding, but touching a wider range of farm activities.

Streeter's The Fat of the Land

The Story of an American Farm, By John Williams Streeter, 12mo, Cloth, 417 pages. \$1.50 net.

Weed's Life Histories of American Insects

By Clarence Moores Weed. 12mo. Cloth. Illus. 248 pages. \$1.50 net.

In these pages are described by an enthusiastic student of entomology such changes as may often be seen in insect form, and which mark the progress of its life.

White's Natural History of Selborne

By Gilbert White. Edited by C. Weeks. 16mo. Cloth. 388 pages, 50 cents; leather, 75 cents.

Wright's Birdcraft

ŧ

A Field Book of Two Hundred Song, Game, and Water Birds. By Mabel Osgood Wright. With Eighty Full-page Plates by Louis Agassis Fuertes, 12mo. Cloth. 339 pages. \$2.00 net.

"The author is not a hard-hearted scientist who goes forth with bag and gun to take life and rob nests, but a patient and intelligent observer who loves the children of the air, and joins their fraternity."—New York Observer.

Wright's Flowers and Ferns in Their Haunts

By Mabel Osgood Wright. With Illustrations from Photographs by the Author and J. Horace McFarland. 12mo, Cloth. 377 pages. \$2.50 net.

"I did not come to pull to pieces or transplant," says the author in the first chapter of this most delightful of outdoor books.

Wise's Diomed

The Life, Travels, and Observations of a Dog. By John Sergeant Wise, Illustrated by J. Linton Chapman. 12mo. Cloth. 377 pages. \$1.50 net.

"One of the best dog stories we know, and the pictures are beautiful, rare for their delicacy and expressiveness, and for their truthfulness to the personality and environment of the dog." — The Literary World.

Professor Lowell's Notable Study of Mars

Mars as the Abode of Life

The book is based on a course of lectures delivered at the Lowell Institute in 1906, supplemented by the results of later observations. It is, in the large, the presentation of the results of the author's research into the genesis and development of what we call a world; not the mere aggregating of matter, but the process by which that matter comes to be individual as we find it. He bridges with the new science of planetology the evolutionary gap between the nebular hypothesis and the Darwinian theory.

It was during the delivery of this series of Lowell Lectures that the Listener in the *Boston Transcript* described his reasoning as "racy, brilliant in style, daring in imagination, building a convincing thesis."

By the Same Author

Previously Published

Mars and Its Canals

"The book makes fascinating reading, and is intended for the average man of intelligence and scientific curiosity. It represents mature reflection, patient investigation and observation, and eleven years' additional work and verification. . . . It is the work of a scientist who has found inspiration and joy in his work; it is full of enthusiasm, but the enthusiasm is not allowed to influence unduly a single conclusion. . . . The book is full of quotable sayings and delightful things. . . . We have here science, reality, that is infinitely stranger and more thrilling than the most imaginative fiction." — Chicago Evening Post.

Each is illustrated with reproductions of exceptional photographs, some of them in colored plates.

Price of each, \$2.50 net; by mail, \$2.70.

PUBLISHED BY

THE MACMILLAN COMPANY

64-66 Fifth Avenue, New York

ivered at the sults of later of the results elopment of matter, but idual as we tology the d the Dar-

ures that asoning ilding a

blished

r the epreion, !t is his red

ull re

ıg

THIS BOOK IS DUE ON THE LAST DATE STAMPED BELOW

AN INITIAL FINE OF 25 CENTS

WILL BE ASSESSED FOR FAILURE TO RETURN THIS BOOK ON THE DATE DUE. THE PENALTY WILL INCREASE TO 50 CENTS ON THE FOURTH DAY AND TO \$1.00 ON THE SEVENTH DAY OVERDUE.

DEC 1 1933	17Feb65\$G
OCT 15 1940	REC'D LD
MAY 26 1941 M	
DEC 2 1941U	
18May'49 W	
18Jul5111y	
Silar 54 IEI	
MAR 18 1954 LU	
	LD 21-100m-7,'33

YB 17001 QB801 Milham. 195096

